

Публикации оппонента Зайцева Александра Михайловича

1. G. Aad *et al.* [ATLAS Collaboration], “Search for the X_b and other hidden-beauty states in the $\pi^+\pi^-\Upsilon(1S)$ channel at ATLAS,” Phys. Lett. B **740**, 199 (2015) [[arXiv:1410.4409 \[hep-ex\]](#)].
2. G. Aad *et al.* [ATLAS Collaboration], “Observation of an Excited B_c^\pm Meson State with the ATLAS Detector,” Phys. Rev. Lett. **113**, no. 21, 212004 (2014) [[arXiv:1407.1032 \[hep-ex\]](#)].
3. G. Aad *et al.* [ATLAS Collaboration], “Measurement of χ_{c1} and χ_{c2} production with $\sqrt{s} = 7$ TeV pp collisions at ATLAS,” JHEP **1407**, 154 (2014) [[arXiv:1404.7035 \[hep-ex\]](#)].
4. G. Aad *et al.* [ATLAS Collaboration], “Measurements of Higgs boson production and couplings in diboson final states with the ATLAS detector at the LHC,” Phys. Lett. B **726**, 88 (2013) [Phys. Lett. B **734**, 406 (2014)] [[arXiv:1307.1427 \[hep-ex\]](#)].
5. G. Aad *et al.* [ATLAS Collaboration], “Measurement of Upsilon production in 7 TeV pp collisions at ATLAS,” Phys. Rev. D **87**, no. 5, 052004 (2013) [[arXiv:1211.7255 \[hep-ex\]](#)].
6. G. Aad *et al.* [ATLAS Collaboration], “Evidence for the spin-0 nature of the Higgs boson using ATLAS data,” Phys. Lett. B **726**, 120 (2013) [[arXiv:1307.1432 \[hep-ex\]](#)].
7. G. Aad *et al.* [ATLAS Collaboration], “Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC,” Phys. Lett. B **716**, 1 (2012) [[arXiv:1207.7214 \[hep-ex\]](#)].
8. G. Aad *et al.* [ATLAS Collaboration], “Time-dependent angular analysis of the decay $B_s^0 \rightarrow J/\psi\phi$ and extraction of $\Delta\Gamma_s$ and the CP-violating weak phase ϕ_s by ATLAS,” JHEP **1212**, 072 (2012) [[arXiv:1208.0572 \[hep-ex\]](#)].
9. G. Aad *et al.* [ATLAS Collaboration], “Observation of a new χ_b state in radiative transitions to $\Upsilon(1S)$ and $\Upsilon(2S)$ at ATLAS,” Phys. Rev. Lett. **108**, 152001 (2012) [[arXiv:1112.5154 \[hep-ex\]](#)].
10. G. Aad *et al.* [ATLAS Collaboration], “Search for squarks and gluinos using final states with jets and missing transverse momentum with the ATLAS detector in $\sqrt{s} = 7$ TeV proton-proton collisions,” Phys. Lett. B **710**, 67 (2012) [[arXiv:1109.6572 \[hep-ex\]](#)].
11. G. Aad *et al.* [ATLAS Collaboration], “Combined search for the Standard Model Higgs boson using up to 4.9 fb^{-1} of pp collision data at $\sqrt{s} = 7$ TeV with the ATLAS detector at the LHC,” Phys. Lett. B **710**, 49 (2012) [[arXiv:1202.1408 \[hep-ex\]](#)].
12. G. Aad *et al.* [ATLAS Collaboration], “Measurement of the top quark-pair production cross section with ATLAS in pp collisions at $\sqrt{s} = 7$ TeV,” Eur. Phys. J. C **71**, 1577 (2011) [[arXiv:1012.1792 \[hep-ex\]](#)].